

УДК 517.54

*I. V. Denega*¹, *A. L. Targonskii*²

¹(*Institute of Mathematics of the National Academy of Sciences of Ukraine, Kyiv*)

²(*Zhytomyr Ivan Franko State University, Zhytomyr*)

¹ iradenega@yandex.ru, ² targonsk@zu.edu.ua

Separating transformation in a problem on extremal decomposition of the complex plane

Dedicated to Prof. Yu. B. Zelinskii on the occasion of his 70th birthday

The paper is devoted to investigation of the problems of geometric function theory of a complex variable. A general problem of the description of extremal configurations maximizing the product of the inner radii of mutually non-overlapping domains is studied.

Робота присвячена дослідженню відкритих проблем геометричної теорії функцій комплексної змінної. Зокрема, вивчається загальна проблема з описання екстремальних конфігурацій, що мінімізують добуток внутрішніх радіусів попарно неперетинних областей.

Let \mathbb{N} and \mathbb{R} be the sets of natural and real numbers, respectively; \mathbb{C} be the Complex plane and let $\overline{\mathbb{C}} = \mathbb{C} \cup \{\infty\}$ be its one-point compactification, $\mathbb{R}^+ = (0, \infty)$. Let $r(B, a)$ be the inner radius of the domain $B \subset \overline{\mathbb{C}}$ with respect to a point $a \in B$ [1, 2]. An inner radius is a generalization of a conformal radius for multiply connected domains.

Definition 1. *A finite collection of arbitrary fixed domains $\{B_k\}_{k=1}^n$, $n \in \mathbb{N}$, $n \geq 2$, such as $B_k \subset \overline{\mathbb{C}}$, $B_k \cap B_m = \emptyset$, $k \neq m$, $k, m = \overline{1, n}$ is called a system of non-overlapping domains.*

Definition 2. *Let $n \in \mathbb{N}$, $n \geq 2$. A set of points $A_n := \{a_k \in \mathbb{C} : k = \overline{1, n}\}$ is called n -radial system if: $|a_k| \in \mathbb{R}^+$, $k = \overline{1, n}$, $0 = \arg a_1 < \arg a_2 < \dots < \arg a_n < 2\pi$.*

Denote $P_k(A_n) := \{w : \arg a_k < \arg w < \arg a_{k+1}\}$, $\theta_k := \arg a_k, a_{n+1} := a_1, \theta_{n+1} := 2\pi, \alpha_k := \frac{1}{\pi} \arg \frac{a_{k+1}}{a_k}, \alpha_{n+1} := \alpha_1, k = \overline{1, n}$. Let $\chi(t) = \frac{1}{2}(t + t^{-1})$. For an arbitrary n -radial system of points $A_n = \{a_k\}, k = \overline{1, n}$, we assume

$$\mathcal{L}(A_n) := \prod_{k=1}^n \chi \left(\left| \frac{a_k}{a_{k+1}} \right|^{\frac{1}{2\alpha_k}} \right) |a_k|.$$

The class of n -radial systems of points for which $\mathcal{L}(A_n) = 1$ automatically includes all systems of n distinct points that are located on the unit circle.

The goal of the present work is the construction of sharp upper bounds for a functional of the form

$$J_n(\gamma) = [r(B_0, 0) r(B_\infty, \infty)]^\gamma \prod_{k=1}^n r(B_k, a_k),$$

where $\gamma \in \mathbb{R}^+, A_n = \{a_k\}_{k=1}^n$ is n -radial system of points, $B_0, B_\infty, \{B_k\}_{k=1}^n$ is system of pairwise disjoint domains, $a_k \in B_k \subset \overline{\mathbb{C}}, k = \overline{0, n}, \infty \in B_\infty \subset \overline{\mathbb{C}}$.

Note that to describe the extremal configurations of the domains we use notion of quadratic differential (see, for example, [1, P. 63–70]).

The following statement holds

Theorem 1. *Let $n \in \mathbb{N}, n \geq 7, 0 < \gamma \leq \gamma_n, \gamma_n = 0,08 n^2$. Then for any n -radial system of points $A_n = \{a_k\}_{k=1}^n$ such that $\mathcal{L}(A_n) = 1$ and any set of mutually non-overlapping domains $B_k, a_0 = 0 \in B_0 \subset \overline{\mathbb{C}}, \infty \in B_\infty \subset \overline{\mathbb{C}}, a_k \in B_k \subset \overline{\mathbb{C}}, k = \overline{1, n}$, the following inequality holds*

$$[r(B_0, 0) r(B_\infty, \infty)]^\gamma \prod_{k=1}^n r(B_k, a_k) \leq [r(\Lambda_0, 0) r(\Lambda_\infty, \infty)]^\gamma \prod_{k=1}^n r(\Lambda_k, \lambda_k),$$

where domains $\Lambda_0, \Lambda_\infty, \Lambda_k$, and points $0, \infty, \lambda_k, k = \overline{1, n}$, are, respectively, circular domains and poles of the quadratic differential

$$Q(w)dw^2 = -\frac{\gamma w^{2n} + (n^2 - 2\gamma)w^n + \gamma}{w^2(w^n - 1)^2} dw^2. \tag{1}$$

Proof. Consider the system of functions

$$\zeta = \pi_k(w) = -i \left(e^{-i\theta_k w} \right)^{\frac{1}{\alpha_k}}, \quad k = \overline{1, n}.$$

The family of functions $\{\pi_k(w)\}_{k=1}^n$ is called admissible for separating transformation of domains $B_0, B_\infty, B_k, k = \overline{1, n}$, with respect to the angles $\{P_k\}_{k=1}^n$. Let $\Omega_k^{(1)}, k = \overline{1, n}$, denote a domain of the plane \mathbb{C}_ζ obtained as a result of the union of the connected component of the set $\pi_k(B_k \cap \overline{P}_k)$ containing the point $\pi_k(a_k)$ with the own symmetric reflection relative to the imaginary axis. In turn, by $\Omega_k^{(2)}, k = \overline{1, n}$, we denote the domain of the plane \mathbb{C}_ζ obtained as a result of the union of the connected component of the set $\pi_k(B_{k+1} \cap \overline{P}_k)$ containing the point $\pi_k(a_{k+1})$ with the own symmetric reflection relative to the imaginary axis, $B_{n+1} := B_1, \pi_n(a_{n+1}) := \pi_n(a_1)$. In addition, by $\Omega_k^{(0)}$ we denote the domain of the plane \mathbb{C}_ζ obtained as a result of the union of the connected component of the set $\pi_k(B_0 \cap \overline{P}_k)$ containing the point $\zeta = 0$ with the own symmetric reflection relative to the imaginary axis. Accordingly, by $\Omega_k^{(\infty)}$ we denote the domain of the plane \mathbb{C}_ζ obtained as a result of the union of the connected component of the set $\pi_k(B_\infty \cap \overline{P}_k)$ containing the point $\zeta = \infty$ with the own symmetric reflection relative to the imaginary axis. Denote $\pi_k(a_k) := \omega_k^{(1)}, \pi_k(a_{k+1}) := \omega_k^{(2)}, k = \overline{1, n}$. It follows from the definition of the function $\pi_k(w)$ that

$$|\pi_k(w) - \omega_k^{(1)}| \sim \frac{1}{\alpha_k} |a_k|^{\frac{1}{\alpha_k} - 1} \cdot |w - a_k|, \quad w \rightarrow a_k, \quad w \in \overline{P}_k,$$

$$|\pi_k(w) - \omega_k^{(2)}| \sim \frac{1}{\alpha_k} |a_{k+1}|^{\frac{1}{\alpha_k} - 1} \cdot |w - a_{k+1}|, \quad w \rightarrow a_{k+1}, \quad w \in \overline{P}_k,$$

$$|\pi_k(w)| \sim |w|^{\frac{1}{\alpha_k}}, \quad w \rightarrow 0, \quad w \in \overline{P}_k,$$

$$|\pi_k(w)| \sim |w|^{\frac{1}{\alpha_k}}, \quad w \rightarrow \infty, \quad w \in \overline{P}_k.$$

Taking into account corresponding results of papers [2, 3], we have the inequalities

$$r(B_k, a_k) \leq \left[\frac{r(\Omega_k^{(1)}, \omega_k^{(1)}) \cdot r(\Omega_k^{(2)}, \omega_k^{(2)})}{\frac{1}{\alpha_k} |a_k|^{\frac{1}{\alpha_k} - 1} \cdot \frac{1}{\alpha_{k-1}} |a_k|^{\frac{1}{\alpha_{k-1}} - 1}} \right]^{\frac{1}{2}}, \quad k = \overline{1, n}, \quad (2)$$

$$\begin{aligned}
r(B_0, 0) &\leq \left[\prod_{k=1}^n r^{\alpha_k^2} \left(\Omega_k^{(0)}, 0 \right) \right]^{\frac{1}{2}}, \\
r(B_\infty, \infty) &\leq \left[\prod_{k=1}^n r^{\alpha_k^2} \left(\Omega_k^{(\infty)}, \infty \right) \right]^{\frac{1}{2}}. \tag{3}
\end{aligned}$$

An equality in (2)–(3) is fully investigated in [2, Theorem 1.9]. By using this inequalities we have the expansion

$$\begin{aligned}
J_n(\gamma) &\leq \prod_{k=1}^n \left(r \left(\Omega_k^{(0)}, 0 \right) r \left(\Omega_k^{(\infty)}, \infty \right) \right)^{\frac{\gamma \alpha_k^2}{2}} \times \\
&\times \left(\frac{r \left(\Omega_k^{(1)}, \omega_k^{(1)} \right) \cdot r \left(\Omega_k^{(2)}, \omega_k^{(2)} \right)}{\frac{1}{\alpha_k} |a_k|^{\frac{1}{\alpha_k} - 1} \cdot \frac{1}{\alpha_{k-1}} |a_k|^{\frac{1}{\alpha_{k-1}} - 1}} \right)^{\frac{1}{2}}.
\end{aligned}$$

Elementary calculations show that

$$\begin{aligned}
J_n(\gamma) &\leq \left(\prod_{k=1}^n \alpha_k \right) \prod_{k=1}^n \frac{|a_k|^{\frac{1}{\alpha_k}} + |a_{k+1}|^{\frac{1}{\alpha_k}}}{(|a_k| |a_{k+1}|)^{\frac{1}{2\alpha_k}}} \cdot |a_k| \times \\
&\times \left\{ \prod_{k=1}^n \left(r \left(\Omega_k^{(0)}, 0 \right) r \left(\Omega_k^{(\infty)}, \infty \right) \right)^{\gamma \alpha_k^2} \cdot \frac{r \left(\Omega_k^{(1)}, \omega_k^{(1)} \right) \cdot r \left(\Omega_k^{(2)}, \omega_k^{(2)} \right)}{\left(|a_k|^{\frac{1}{\alpha_k}} + |a_{k+1}|^{\frac{1}{\alpha_k}} \right)^2} \right\}^{\frac{1}{2}}.
\end{aligned}$$

Hence

$$\begin{aligned}
J_n(\gamma) &\leq \left(\prod_{k=1}^n \alpha_k \right) \prod_{k=1}^n \left(\left| \frac{a_k}{a_{k+1}} \right|^{\frac{1}{2\alpha_k}} + \left| \frac{a_{k+1}}{a_k} \right|^{\frac{1}{2\alpha_k}} \right) |a_k| \times \\
&\times \left\{ \prod_{k=1}^n \left(r \left(\Omega_k^{(0)}, 0 \right) r \left(\Omega_k^{(\infty)}, \infty \right) \right)^{\gamma \alpha_k^2} \cdot \frac{r \left(\Omega_k^{(1)}, \omega_k^{(1)} \right) \cdot r \left(\Omega_k^{(2)}, \omega_k^{(2)} \right)}{\left(|a_k|^{\frac{1}{\alpha_k}} + |a_{k+1}|^{\frac{1}{\alpha_k}} \right)^2} \right\}^{\frac{1}{2}}.
\end{aligned}$$

Thus

$$\prod_{k=1}^n \left(\left| \frac{a_k}{a_{k+1}} \right|^{\frac{1}{2\alpha_k}} + \left| \frac{a_{k+1}}{a_k} \right|^{\frac{1}{2\alpha_k}} \right) |a_k| = \prod_{k=1}^n \chi \left(\left| \frac{a_k}{a_{k+1}} \right|^{\frac{1}{2\alpha_k}} \right) |a_k| = \mathcal{L}(A_n),$$

and we obtain

$$J_n(\gamma) \leq 2^n \cdot \left(\prod_{k=1}^n \alpha_k \right) \cdot \mathcal{L}(A_n) \times \\ \times \prod_{k=1}^n \left\{ \frac{r(\Omega_k^{(1)}, \omega_k^{(1)}) \cdot r(\Omega_k^{(2)}, \omega_k^{(2)})}{\left(|a_k|^{\frac{1}{\alpha_k}} + |a_{k+1}|^{\frac{1}{\alpha_k}} \right)^2} \left(r(\Omega_k^{(0)}, 0) r(\Omega_k^{(\infty)}, \infty) \right)^{\gamma \alpha_k^2} \right\}^{\frac{1}{2}}.$$

For convenience, right side of the last inequality we multiply and divide by the value $\sqrt{\gamma}$ and get the inequality

$$J_n(\gamma) \leq \left(\frac{2}{\sqrt{\gamma}} \right)^n \cdot \left(\prod_{k=1}^n \alpha_k \sqrt{\gamma} \right) \cdot \mathcal{L}(A_n) \times \\ \times \prod_{k=1}^n \left\{ \frac{r(\Omega_k^{(1)}, \omega_k^{(1)}) \cdot r(\Omega_k^{(2)}, \omega_k^{(2)})}{\left(|a_k|^{\frac{1}{\alpha_k}} + |a_{k+1}|^{\frac{1}{\alpha_k}} \right)^2} \left(r(\Omega_k^{(0)}, 0) r(\Omega_k^{(\infty)}, \infty) \right)^{\gamma \alpha_k^2} \right\}^{\frac{1}{2}},$$

where $|a_k|^{\frac{1}{\alpha_k}} + |a_{k+1}|^{\frac{1}{\alpha_k}} = |\omega_k^{(2)} - \omega_k^{(1)}|$, $k = \overline{1, n}$. Equality in the last inequality is attained if and only if the equality is attained in inequalities (2) – (3) for all $k = \overline{1, n}$. Each expression contained in braces in the last inequality is value of the functional

$$K_\tau = [r(B_0, 0) r(B_\infty, \infty)]^{\tau^2} \cdot \frac{r(B_1, a_1) r(B_2, a_2)}{|a_1 - a_2|^2} \quad (4)$$

on the system of non-overlapping domains $\{\Omega_k^{(0)}, \Omega_k^{(1)}, \Omega_k^{(2)}, \Omega_k^{(\infty)}\}$ and corresponding points system $\{0, \omega_k^{(1)}, \omega_k^{(2)}, \infty\}$, $k = \overline{1, n}$. Functional evaluation (4) in the case of fixed poles was first obtained by V. N. Dubinin, then by G. V. Kuzmina, E. Emelyanov, A. L. Tarhonskii. Note that in our case the points a_1 and a_2 of the functional (4) are not fixed and not symmetrical. So, fractional-linear transformations can not reduce them to -1 and 1 . V. N. Dubinin and G. V. Kuzmina considered the case when

the points a_1 and a_2 are symmetrical and equal to 1 and -1 , respectively. Therefore decisive role is played corollary of Theorem 4.1.1 in [1, p. 169]. From here, based on invariance of the functional (4) as in the proof of Theorem 4.1.1 in [1], we have that $K_\tau \leq \Phi(\tau)$, $\tau \geq 0$, where $\Phi(\tau) = \tau^{2\tau^2} \cdot |1 - \tau|^{-(1-\tau)^2} \cdot (1 + \tau)^{-(1+\tau)^2}$. Then

$$\begin{aligned} J_n(\gamma) &\leq \left(\frac{2}{\sqrt{\gamma}}\right)^n \left(\prod_{k=1}^n \alpha_k \sqrt{\gamma}\right) \left[\prod_{k=1}^n \Phi(\tau_k)\right]^{1/2} = \\ &= \left(\frac{2}{\sqrt{\gamma}}\right)^n \left[\prod_{k=1}^n \left(\tau_k^{2\tau_k^2+2} \cdot |1 - \tau_k|^{-(1-\tau_k)^2} \cdot (1 + \tau_k)^{-(1+\tau_k)^2}\right)\right]^{\frac{1}{2}}, \end{aligned}$$

where $\tau_k = \sqrt{\gamma} \cdot \alpha_k$, $k = \overline{1, n}$. Let

$$S(x) = x^{2x^2+2} \cdot |1 - x|^{-(1-x)^2} \cdot (1 + x)^{-(1+x)^2} \quad \text{and} \quad \Psi(x) = \ln(S(x)).$$

$S(x)$ is logarithmically convex function on the interval $[0, x_0]$, $x_0 \approx 0,88441$. Further similarly [3, 4] we consider the following extremal problem

$$\prod_{k=1}^n S(x_k) \longrightarrow \max, \quad \sum_{k=1}^n x_k = 2\sqrt{\gamma}, \quad x_k = \alpha_k \sqrt{\gamma}.$$

Let $X^{(0)} = \{x_k^{(0)}\}_{k=1}^n$ be any extremal set of points in above mentioned problem. In a similar way as in [4], we obtain the following result: if $0 < x_k^{(0)} < x_j^{(0)}$, then

$$\Psi'(x_k^{(0)}) = \Psi'(x_j^{(0)}), \quad (5)$$

where $k, j = \overline{1, n}$, $k \neq j$,

$$\Psi'(x) = 4x \ln(x) - 2(x-1) \ln|x-1| - 2(x+1) \ln(x+1) + \frac{2}{x}$$

(see Fig. 1).




Fig. 1: Graph of the function $y = \Psi'(x)$

By using similar arguments as in the paper [4] and the relation (5) we will prove that $x_1^{(0)} = x_2^{(0)} = \dots = x_n^{(0)}$. Taking into account properties of the function $\Psi'(x)$ and conditions of Theorem we obtain the following relation: $(x_1 - 0,56)n + (x_2 - x_1) > 0$ for $n \geq 7$. Therefore, we have $nx_1 + (x_2 - x_1) > 0,56n$. Finally we get

$$(n-1)x_1 + x_2 > 2\sqrt{\gamma_n}, \quad \gamma_n = 0,08n^2, \quad n \geq 7.$$

Thus, in accordance with [4], we agree to say that the set of points $\{x_k^{(0)}\}_{k=1}^n$, $n \geq 7$, can not be extremal if $x_n^{(0)} \in (x_0, 2]$. Consequently, for an extremal set $\{x_k^{(0)}\}_{k=1}^n$ is only possible in the case when $x_k^{(0)} \in (0, x_0]$, $k = \overline{1, n}$, and $x_1^{(0)} = x_2^{(0)} = \dots = x_n^{(0)}$. For $\gamma < \gamma_n$, $n \geq 7$, all previ-

ous arguments are valid. The equality case is straightforward to verify. Theorem 1 is proved.

Corollary 1. *Under conditions of Theorem 1 the following inequality holds*

$$[r(B_0, 0)r(B_\infty, \infty)]^\gamma \prod_{k=1}^n r(B_k, a_k) \leq \left(\frac{4}{n} \right)^n \left[\frac{\left(\frac{4\gamma}{n^2} \right)^{\left(\frac{4\gamma}{n^2} \right)}}{\left| \frac{2\sqrt{\gamma}}{n} - 1 \right|^{\left(\frac{2\sqrt{\gamma}}{n} - 1 \right)^2} \left(\frac{2\sqrt{\gamma}}{n} + 1 \right)^{\left(\frac{2\sqrt{\gamma}}{n} + 1 \right)^2}} \right]^{\frac{n}{2}}.$$

Equality is attained if $0, \infty, a_k$, and $B_0, B_\infty, B_k, k = \overline{1, n}$, are, respectively, poles and circular domains of the quadratic differential (1).

Corollary 2. *Let $n \in \mathbb{N}, n \geq 7, 0 < \gamma \leq \gamma_n, \gamma_n = 0,08n^2$. Then for any different points on the circle $|a_k| = 1, k = \overline{1, n}$, and any pairwise non-overlapping domains $B_k, a_0 = 0 \in B_0 \subset \overline{\mathbb{C}}, \infty \in B_\infty \subset \overline{\mathbb{C}}, a_k \in B_k \subset \overline{\mathbb{C}}, k = \overline{1, n}$, the following inequality holds*

$$[r(B_0, 0)r(B_\infty, \infty)]^\gamma \prod_{k=1}^n r(B_k, a_k) \leq [r(\Lambda_0, 0)r(\Lambda_\infty, \infty)]^\gamma \prod_{k=1}^n r(\Lambda_k, \lambda_k),$$

where domains $\Lambda_0, \Lambda_\infty, \Lambda_k$, and points $0, \infty, \lambda_k, k = \overline{1, n}$, are, respectively, circular domains and poles of the quadratic differential (1).

References

- [1] *Bakhtin A.K., Bakhtina G.P., Zelinskii Yu.B.* Topological-algebraic structures and geometric methods in complex analysis // Proceedings of the Institute of Mathematics of NAS of Ukraine, 2008. — 308 p. (in Russian).
- [2] *Dubinina V.N.* Symmetrization method in geometric function theory of complex variables // Successes Mat. Science. — 1994. — **49**, No. 1 (295). — P. 3 – 76 (in Russian); Engl. transl. in: Russian Math. Surveys. — 1994. — **49**, № 1. — P. 1 – 79.

- [3] *Dubinin V. N.* Separating transformation of domains and problems on extremal decomposition // Notes scientific. sem. Leningr. Dep. of Math. Inst. AN USSR. — 1988. — **168**. — P. 48 – 66 (in Russian); Engl. transl. in: J. Soviet Math. — 1991. — **53**, No. 3. — P. 252 – 263.
- [4] *Kovalev L. V.* On the problem of extremal decomposition with free poles on a circle // Dal'nevostochnyi Mat. Sb. — 1996. — **2**. — P. 96 – 98 (in Russian).
- [5] *Lavrentev M. A.* To the theory of conformal mappings // Trudy Fiz.-Mat. Inst. AN SSSR. — 1934. — **5**. — P. 159 – 245 (in Russian).
- [6] *Goluzin G. M.* Geometric theory of functions of a complex variable. — Amer. Math. Soc., Providence, R.I., 1969. — 676 p.
- [7] *Hayman V. K.* Multivalent functions. — Cambridge University Press, 1958. — 151 p.
- [8] *Jenkins J. A.* Univalent functions and conformal mapping. — M.: Publishing House of Foreign Literature, 1962. — 256 p. (in Russian).
- [9] *Kolbina L. I.* Conformal mapping of the unit circle onto non-overlapping domains // Bulletin of Leningrad University. — 1955. — **5**. — P. 37 – 43 (in Russian).
- [10] *Kuzmina G. V.* Problems on extremal decomposition of the riemann sphere // Notes scientific. sem. Leningr. Dep. of Math. Inst. AN USSR. — 2001. — **276**. — P. 253 – 275 (in Russian); Engl. transl. in: J. of Math. Sci — 2003. — **118**, No. 1. — P. 4880 – 4894.
- [11] *Bakhtin A. K., Denega I. V.* Addendum to a theorem on extremal decomposition of the complex plane // Bulletin de la société des sciences et des lettres de Łódź, Recherches sur les déformations. — 2012. — **62**, № 2. — P. 83 – 92.
- [12] *Bakhtin A. K., Bakhtina G. P., Vjun V. E.* On some inequalities in the theory of non-overlapping domains // Zb. Pr. Inst. Mat. NAN Ukr. — K.: In-te of mathematics of NAS of Ukraine, 2014. — **11**, № 1. — P. 141 – 152 (Ukrainian).
- [13] *Dvorak I. Ya.* On the inner radii of symmetric nonoverlapping domains // Ukrainskii Matematychnyi Visnyk. — 2016. — **13**, No. 2. — P. 10 – 19 (in Russian).